

-56-

## CLAIMS

We claim:

- 5
1. A humanized CC49 antibody, comprising a non-conservative amino acid substitution in a light chain complementarity determining region 3 of the CC49 antibody, or functional fragment of the humanized CC49 antibody, that has a high binding affinity for TAG-72.
  - 10 2. The antibody of claim 1, wherein the non-conservative substitution is a tyrosine to proline substitution.
  3. The antibody of claim 1, wherein the non-conservative substitution is at position  
15 91.
  4. The antibody of claim 1, wherein the non-conservative substitution is at a residue that is a ligand contact residue.
  - 20 5. The antibody of claim 1, wherein the functional fragment is an Fab fragment, an Fv fragment, or an F(ab')<sub>2</sub> fragment.
  6. The antibody of claim 1, wherein a light chain complementarity determining region 1 and a light chain complementarity determining region 2 are from a human  
25 antibody.
  7. The antibody of claim 1, wherein the light chain complementarity determining region 3, a heavy chain complementarity determining region 1, a heavy chain

-57-

complementarity determining region 2, and a heavy chain complementarity determining region 3 are from a murine CC49 antibody.

- 5      8.      The antibody of claim 1, wherein the high binding affinity is at least about  $1.2 \times 10^{-8}$  M.
9.      The antibody of claim 8, wherein the high binding affinity is at least about  $1.5 \times 10^{-8}$ , about  $2.0 \times 10^{-8}$ , about  $2.5 \times 10^{-8}$ , about  $3.0 \times 10^{-8}$ , about  $3.5 \times 10^{-8}$ , about  $4.0 \times 10^{-8}$ , about  $4.5 \times 10^{-8}$ , or about  $5.0 \times 10^{-8}$  M.
- 10      10.      The antibody of claim 1, wherein the antibody is minimally immunogenic.
11.      The antibody of claim 1, wherein the antibody further comprises an effector molecule.
- 15      12.      The antibody of claim 11, wherein the effector molecule is a detectable label.
13.      The antibody of claim 12, wherein the detectable label comprises a radioactive isotope, an enzyme substrate, a co-factor, a ligand, a chemiluminescent agent, a fluorescent agent, a hapten, or an enzyme.
- 20      14.      The antibody of claim 11, wherein the effector molecule is a toxin.
15.      The antibody of claim 14, wherein the toxin is a chemotherapeutic drug, a radioactive isotope, a bacterial toxin, a viral toxin, a cytokine or a venom protein.
- 25      16.      The antibody of claim 1, further comprising at least one additional non-conservative amino acid substitution in a light chain complementarity determining region.

-58-

17. The antibody of claim 16, wherein the light chain complementarity determining region is a light chain complementarity determining region 1, a light chain complementarity determining region 2, or a light chain complementarity determining region 3.

18. The antibody of claim 1, further comprising at least one non-conservative amino acid substitution in a heavy chain complementarity determining region.

19. The antibody of claim 18, wherein the heavy chain complementarity determining region is a heavy chain complementarity determining region 1, a heavy chain complementarity determining region 2, or a heavy chain complementarity determining region 3.

20. A humanized CC49 antibody, wherein a nucleic acid sequence encoding the antibody has an ATCC Accession number comprising ATCC Accession number PTA-4182 or ATCC Accession number PTA-4183.

21. A nucleic acid molecule encoding the humanized monoclonal antibody of claim 1.

22. A vector comprising the nucleic acid of claim 21.

23. A humanized CC49 antibody, comprising:  
a variable light framework region and a variable heavy framework region of a human antibody;  
a complementarity determining region, wherein at least one complementarity determining region is from the human antibody and the remaining complementarity determining regions are from a murine CC49 antibody;

-59-

a non-conservative substitution of a first residue in a light chain complementarity determining region 3; and

a substitution of a second residue in a complementarity determining region of the human CC49 antibody;

5                wherein the humanized CC49 antibody has a high binding affinity for TAG-72 and is minimally immunogenic.

24.    The antibody of claim 23, wherein the non-conservative substitution is a tyrosine to proline substitution.

10

25.    The antibody of claim 23, wherein the non-conservative substitution is at position 91.

26.    The antibody of claim 25, wherein the non-conservative substitution at position  
15    91 is a tyrosine to proline substitution.

27.    The antibody of claim 23, wherein the antibody further comprises an effector molecule.

20    28.    The antibody of claim 27, wherein the effector molecule is a detectable label.

29.    The antibody of claim 28, wherein the detectable label comprises a radioactive isotope, an enzyme substrate, a co-factor, a ligand, a chemiluminescent agent, a fluorescent agent, a hapten, or an enzyme.

25

30.    The antibody of claim 27, wherein the effector molecule is a toxin.

31.    The antibody of claim 30, wherein the toxin is a chemotherapeutic drug, a radioactive isotope, a bacterial toxin, a viral toxin, a cytokine or a venom protein.

-60-

32. A method of detecting a TAG-72-expressing tumor in a subject, comprising:  
contacting a sample obtained from the subject with the antibody of claim  
1 for a sufficient amount of time to form an immune complex;  
5 detecting the presence of the immune complex, wherein the presence of  
the immune complex demonstrates the presence of the TAG-72-expressing tumor.
33. The method of claim 32, wherein the tumor is a colorectal tumor, a gastric  
tumor, a pancreatic tumor, a breast tumor, a lung tumor, an adenocarcinoma, or an  
10 ovarian tumor.
34. The method of claim 32, wherein the antibody further comprises an effector  
molecule.
- 15 35. The method of claim 34, wherein the effector molecule is a detectable label.
36. The method of claim 35, wherein the detectable label comprises a radioactive  
isotope, an enzyme substrate, a co-factor, a ligand, a chemiluminescent agent, a  
fluorescent agent, a hapten, or an enzyme.  
20
37. The method of claim 32, further comprising contacting the antibody with a  
secondary antibody.
38. The method of claim 37, wherein the secondary antibody further comprises a  
25 detectable label.
39. A method of detecting a TAG-72-expressing tumor in a subject, comprising:  
administering the antibody of claim 1 to the subject for a sufficient  
amount of time to form an immune complex;

REPL  
ART 34 AM

-61-

detecting the presence of the immune complex, wherein the presence of the immune complex demonstrates the presence of the TAG-72-expressing tumor.

40. The method of claim 39, wherein the antibody further comprises an effector  
5 molecule.

41. The method of claim 40, wherein the effector molecule is a detectable label.

42. The method of claim 41, wherein the detectable label comprises a radioactive  
10 isotope, an enzyme substrate, a co-factor, a ligand, a chemiluminescent agent, a fluorescent agent, a hapten, or an enzyme.

43. The method of claim 39, wherein the tumor is a colorectal tumor, a gastric  
tumor, a pancreatic tumor, a breast tumor, a lung tumor, an adenocarcinoma, or an  
15 ovarian tumor.

44. A method of treating a subject having a tumor that expresses TAG-72,  
comprising administering to the subject a therapeutically effective amount of the  
antibody of claim 1, wherein administering the therapeutically effective amount of the  
20 antibody of claim 1 inhibits the growth of the tumor or reduces the size of the tumor,  
thereby treating the subject.

45. The method of claim 44, wherein the administration of a therapeutically  
effective amount of the antibody of claim 1 does not elicit a human anti-murine  
25 antibody response in a subject.

46. The method of claim 44, wherein the tumor is a colorectal tumor, a gastric  
tumor, a pancreatic tumor, a breast tumor, a lung tumor, an adenocarcinoma, or an  
ovarian tumor.

-62-

47. The method of claim 44, wherein the antibody further comprises an effector molecule.

5 48. The method of claim 47, wherein the effector molecule is a toxin.

49. The method of claim 48, wherein the toxin is a chemotherapeutic drug, a radioactive isotope, a bacterial toxin, a viral toxin, a cytokine, or a venom protein.

10 50. The method of claim 49, wherein the antibody comprising a radioactive isotope is used in radioimmunotherapy.

51. A method of treating a subject having a tumor that expresses TAG-72, comprising:

15 administering the antibody of claim 1 to the subject for a sufficient amount of time to form an immune complex, wherein the antibody comprises a radioactive isotope;

detecting the presence of the immune complex with a hand-held gamma counter, wherein the presence of the immune complex demonstrates the presence of the  
20 TAG-72-expressing tumor; and

removing the tumor surgically, thereby treating the subject.

52. A pharmaceutical composition comprising a therapeutically effective amount of the antibody of claim 1 in a pharmaceutically acceptable carrier.

25

53. A kit, comprising a container comprising the antibody of claim 1.

REPLACED BY  
FIG 24A/B/C

-63-

54. The kit of claim 53, further comprising a container containing an antigen, a container containing a secondary antibody conjugated to a chemical compound, instructions for using the kit, or any combination thereof.